



## Brief Report

## A pilot study of *Trabajadora de salud*, a lay health worker intervention for Latinas/os with traumatic brain injuries and their caregivers



Kristen F. Linton, M.S.W., Ph.D.<sup>a,\*</sup>, Bum Jung Kim, M.S.W., Ph.D.<sup>b</sup>

<sup>a</sup> Health Science Program, California State University, Channel Islands, One University Drive, Solano Hall 1120, Camarillo, CA, 93012, United States

<sup>b</sup> School of Social Work, University of Hawai'i at Manoa, United States

## ARTICLE INFO

## Article history:

Received 13 August 2016

Received in revised form

11 April 2017

Accepted 15 April 2017

## Keywords:

Latino

Hispanic

Brain injury

Traumatic brain injury

Trabajadora

Lay health worker

## ABSTRACT

**Background:** Latinas/os with traumatic brain injuries (TBIs) and their caregivers experience worse outcomes than others.

**Objective:** The study aimed to assess the acceptability and promise of *Trabajadora de Salud* on the functional abilities, hospital readmission, rehabilitation, employment, depression, somatic symptoms, and caregiver burden among Latinas/os with TBIs and their caregivers.

**Methods:** A pre-posttest experimental pilot study was conducted. A total of eight Latina/o adult patients (50% female) with mild or moderate TBI and six of their caregivers (66.7% female) were randomized to receive *Trabajadora de Salud* or a telephone only control group. *Trabajadora de Salud*, a three-month, in-home intervention administered by bilingual lay health workers, focused on: 1) providing empathy and validation of TBI symptoms, 2) addressing basic needs, 3) goal setting, and 4) improving communication with healthcare providers.

**Results:** *Trabajadora de Salud* was widely accepted by patients, caregivers, and health professionals. The functional, depression, and somatic symptoms of the patients as well as the somatic symptoms and caregiver burden of the caregivers improved more for participants in the intervention group than the control group.

**Conclusions:** *Trabajadora de Salud* demonstrated promise in improving outcomes of Latinas/os with TBIs and their caregivers and should be further studied.

© 2017 Elsevier Inc. All rights reserved.

Every year 1.7 million people suffer a traumatic brain injury (TBI) resulting in death or permanent disability.<sup>1</sup> While the incidence rate of the general United States population is 200 per 100,000, it is 262 per 100,000 for the Hispanic population. Latina/o survivors of TBIs are more likely to be males, an average age of 30, and sustain a TBI due to transportation, falls, or violence.<sup>2</sup> Latinas/os are more likely to experience penetrating TBIs, which are associated with significantly higher rates of respiratory failure, pneumonia, cerebrospinal fluid leak, and hypotonia.<sup>3</sup> Latinas/os are 42% more likely to be discharged from inpatient acute care hospitalization to their home versus to intensive rehabilitation (outpatient or inpatient) than whites. This research also found that Latinas/os are less likely to be discharged to a level of rehabilitation consistent with their injury than their matched white counterparts.<sup>4</sup> Latina/o

survivors of TBIs are more likely to sustain long term functional inability, unemployment, and psychological issues than whites.<sup>5,6</sup> Due to the dependence of survivors of TBIs on others, caregivers are essential to supporting the survivor through the emotional recovery process as well as providing essential hands-on support as needed in daily living activities, such as bathing, dressing, and eating, and ensuring that patients attend medical appointments. Additionally, Latina/o caregivers express more burden and spend more time in caregiving roles.<sup>7</sup>

Interventions aimed at improving TBI outcomes among Latinas/os are scarce. The only published intervention solely designed for Latinas/os with TBI included a 10-week group solution-focused brief therapy (SFBT) intervention that concentrated on locus of control, skill acquisition, and family participation.<sup>8</sup> Facilitators were bilingual and bicultural; language, acculturation, migratory stressors, attitudes and beliefs about disability and health care, and support networks were also discussed. Participants had statistically significant improvements on hopelessness ( $t = 7.9, p < 0.005$ ), life

\* Corresponding author.

E-mail address: [Kristen.linton@csuci.edu](mailto:Kristen.linton@csuci.edu) (K.F. Linton).

purpose ( $t = -6.7, p < 0.005$ ), and self-regulatory ability ( $t = 5.6, p < 0.005$ ). Other SFBT psycho-educational interventions have demonstrated efficacy for improving TBI knowledge and psychological adjustment for the general population of people with TBI.<sup>9,10</sup>

The *trabajadora de salud* (lay health worker) model has demonstrated efficacy in improving health care use, psychological, and functional outcomes of Latinas/os and Hispanics with various common health conditions (i.e. diabetes, cancer, mental health).<sup>11</sup> *Trabajadoras*, who are commonly bilingual and bicultural, are trained on the targeted health condition and provide culturally grounded education and/or bridge support for patients. A systematic review of 128 published articles on *trabajadora de salud* interventions for Hispanics found that in all except one study there were observed significant improvements in health behaviors regardless of the health condition targeted.<sup>12</sup>

Section 3025 of the United States Affordable Care Act requires Centers for Medicare and Medicaid to reduce payments to hospitals with excess readmissions.<sup>13</sup> In a recent study of 10,390 individuals hospitalized for a TBI, 35.5% were subsequently hospitalized during a 3-year follow-up period.<sup>14</sup> If *Trabajadora de Salud* is efficacious in improving outcomes among Latinas/os with TBI and their caregivers, this increased knowledge can be used to establish *Trabajadora de Salud* programs to promote equal opportunities for healthy and productive lives among Latinos who have suffered a TBI. This study aimed to assess the acceptability and promise of *Trabajadora de Salud* in improving functional abilities, hospital readmission, rehabilitation, employment, depression, somatic symptoms, and caregiver burden among Latinas/os with TBIs and their caregivers.

## Methods

### Participants

Patient inclusion criteria for participants included: 18 years old or older, self-identification as Latina/o, and had been hospitalized for a TBI caused by a transportation, fall, accident or another external cause within the past year prior to participation in the study. Patient exclusion criteria included: brain injury due to a stroke or internal cause and co-occurring disorders, such as mental illness or physical illness. Caregiver inclusion criteria included: 18 years old or older, identified by the patient as the primary caregiver of the patient, and a family member or friend of the patient. Caregiver exclusion criteria included: provides professional, paid caregiving to the patient. IRB approval was obtained from California State University, Channel Islands prior to recruitment. Patients were recruited from flyers at rehabilitation centers ( $n = 5$ ) and bilingual advertisements in Latina/o newspapers ( $n = 3$ ).

### Measurements

A pre-posttest experimental design was used including measures that were tested for reliability in previous research on Latinas/os with TBIs and their caregivers. All participants were asked questions on demographic information. The following measures were used for patients: the complete Functional Independence Measure (higher scores = more independence) and only the depression and somatic symptom scales of the Neurobehavioral Functioning Inventory (higher scores = more symptoms).<sup>15,16</sup> Patients were also asked if they had received inpatient or outpatient rehabilitation or been hospitalized since their injury. The depression and somatic symptom scales of the Neurobehavioral Functioning Inventory and the complete Caregiver Strain Index (higher scores = more strain) were used for caregivers.<sup>17</sup> Patients and caregivers were both asked if they had been employed full time, part time, or temporarily prior to the patients' injury as well as at

the time of the pre-test and post-test. Patients and caregivers received incentives at pre-test (\$10 gift card) and post-test (\$12 gift card) administration.

### Intervention

Patients and caregivers were randomly assigned to an intervention or control group. The three-month, in-home, *Trabajadora de Salud* intervention was administered by bilingual lay health workers and compared to those in a control group among eight Latina/o adult patients with TBI and six of their caregivers in Ventura, Los Angeles, and Kern Counties, California. The *Trabajadora de Salud* in-person visits utilized SFBT and focused on: 1) providing empathy and validation of TBI symptoms, 2) addressing basic needs, 3) well-formed goal setting especially related to employment, and 4) improving communication with healthcare providers. *Trabajadoras* were trained on SFBT, TBIs, acculturation, migratory stressors, Latina/o attitudes and beliefs toward disability and health care, and community resources for TBI regarding health care and employment. SFBT interventions and other *trabajadora de salud* interventions have typically been brief and empowering; the therapists or *trabajadoras* both teach participants skills that can be used after the intervention is complete.<sup>18,19</sup> *Trabajadora de Salud* followed this brief, empowering model of support. The control group received a less intensive telephone only version of the intervention. The control group received two phone calls over a three month period by a *trabajadora* who focused on the same elements as the *Trabajadora de Salud* in-person intervention. The *trabajadoras* who conducted the telephone only intervention were also bilingual and bicultural and received the same training as the *trabajadoras* who receive the in-person intervention.

*Trabajadora de salud* interventions have demonstrated high retention rates by ensuring that transportation costs were covered and providing telephone reminders prior to each visit by the *trabajadora*. The *trabajadora* met the patient and caregiver at his or her home and conducted reminder phone calls with participants prior to each visit. Telephone reminders increase the development of the relationship between the *trabajadora* and participants thus demonstrating consistency with *personalismo*, a common cultural value among Latinas/os defined as "formal friendliness" and also described as a cultural emphasis on personal relationships.<sup>20</sup>

Due to the small sample size, descriptive statistics only were used to describe and compare outcomes of the intervention and control group.

## Results

Eight patients and six of their corresponding caregivers participated in the study and were randomized into the intervention and control groups. Table 1 includes basic characteristics of the pilot study participants. Three dyads (three patients and three corresponding caregivers) received the full intervention which ranged from 6 to 12 in-person visits ( $M = 8, SD = 3.09$ ), while five patients and two of their corresponding caregivers were in the telephone-only control group. The functional, depression, and somatic symptoms of the patients (Table 2) as well as the somatic symptoms and caregiver burden of the caregivers (Table 3) improved more for participants in the intervention group than the control group as shown by pre-post change in mean score. All patients were hospitalized for their injuries at least once with a range from 1 to 2 times. No patients in the intervention or control group were hospitalized again after the study began.

During the course of the pilot, the acceptability of the intervention was assessed. Patients and health professionals responded positively to the intervention protocol. A caregiver who received

**Table 1**  
Participant demographics.

	Patients (n = 8) M (SD) Range = 24–52	Caregivers (n = 6) M (SD) Range = 32–56
	n (%)	n (%)
Age	36.5 (10.47) Range = 24–52	44.83 (10.55) Range = 32–56
Gender		
Female	4 (50)	5 (66.7)
Self-Identified Ethnicity		
Mexican	7 (87.5)	4 (66.7)
Argentinian	1 (12.5)	1 (16.7)
African American		1 (16.7)
Health Insurance		
MediCal	4 (50.0)	4 (66.7)
Private	2 (25.0)	1 (16.7)
None	2 (25.0)	1 (16.7)
Annual Income		
<sup>a</sup> Less than \$15,654 (133% FPL)	7 (87.5)	5 (80.0)
Education		
Less than High school	2 (25.0)	4 (60.0)
High school graduate	2 (25.0)	2 (40.0)
<sup>a</sup> Some college	2 (25.0)	
College graduate	2 (25.0)	
Language at Home		
English	5 (62.5)	4 (66.7)
Spanish	2 (25.0)	2 (33.4)
Both	1 (12.5)	
Etiology of TBI#1		
Fall	4 (50.0)	
Car Accident	4 (50.0)	

Note.

<sup>a</sup> 133% of the Federal Poverty Level (FPL) was the eligibility criteria for MediCal in the State of California at the time of this study. “Some college” is defined as receipt of college courses with or without credit and without graduation.

*Trabajadora de Salud* voluntarily sent a thank you letter to her *trabajadora* and reported how she saw “measurable improvement in many areas of his [her son’s] disabilities.” She expressed that the *trabajadora*’s “assistance has helped him to move forward rapidly where he was stagnant before. You brought happiness as well as greater communication, social, movement, thinking and cognitive

skills to [my son]”. This powerful voluntary letter describes outcomes that the intervention was aiming to improve. A patient described her *trabajadora* as someone that “helps me a lot and does research for me” referring to the *trabajadora*’s role in finding resources for her.

## Discussion

The pilot study had some notable findings. The study found that the *Trabajadora de Salud* intervention demonstrated promise in improving the functional abilities, depression and somatic symptoms of Latina/o patients with TBIs as well as somatic symptoms and caregiver burden among their caregivers as compared to a less intensive telephone only intervention. Future research is needed to understand the interventions’ promise on improving the patients’ hospital readmission, receipt of rehabilitation, and employment of Latina/o patients with TBIs. A follow-up survey months after the completion of the intervention could improve the likelihood of detecting an effect of the intervention on these outcomes. Depression among the caregivers was measured in the pre and post-tests in this study, yet those in the intervention group did not improve their outcomes more than those in the control group. The caregivers in the sample reported high levels of depression. The brief in-person intervention may not be strong enough to impact severe symptoms of depression.

The caregivers reported higher pretest depression and somatic symptoms than the patients on average, yet no statistical analyses were performed due to the small sample size (Tables 2 and 3). While *Trabajadora de Salud* supports the family unit, the patient was often the focus of the intervention. Caregivers who received the *Trabajadora de Salud* intervention did not improve their depression symptoms more than those in the control group. The intervention protocol for the efficacy study proposed will be adjusted to emphasize the caregiver as a main foci of the intervention to increase the potential of improving their outcomes more.

Secondly, the pilot study found that the intervention group and control group did not differ on receipt on rehabilitation or employment. A barrier to employment is lack of private health

**Table 2**  
Patient outcomes.

	Intervention (n = 3)			Control (n = 5)		
	Pre	Post	Pre-Post Change in Score	Pre	Post	Pre-Post Change in Score
Frequency						
Rehab						
Physical	1	2	1	3	3	0
Occupation	0	0	0	2	2	0
Speech	0	0	0	3	3	0
Employed	0	1	1	2	3	1
M (SD)						
Physical FIM (higher scores = more independence)	62.67 (3.51)	65 (1.00)	2.33	55.40 (19.88)	57 (19.01)	1.6
Cognitive FIM (higher scores = more independence)	22.33 (3.21)	27 (2.00)	4.7	23 (2.46)	23 (4.18)	0
*Depression (higher scores = more depression)	18.67 (5.13)	15.33 (2.51)	−3.34	28.4 (2.96)	26.6 (4.03)	−1.8
*Somatic (higher scores = more somatic symptoms)	17.33 (4.61)	12.67 (1.15)	−4.66	25.2 (3.11)	23.4 (3.36)	−1.8

Note. Depression and Somatic symptoms as measured by the Neurobehavioral Functioning Inventory.

**Table 3**  
Caregiver outcomes.

	Intervention (n = 3)			Control (n = 3)		
	Pre	Post	Pre-Post Change in Score	Pre	Post	Pre-Post Change in Score
Depression (higher scores = more depression)	22.5 (5.56)	18.75 (2.98)	−3.75	30.00 (9.00)	23.67 (5.68)	−6.33
Somatic (higher scores = more somatic symptoms)	28 (11.31)	17 (4.75)	−11	28.33 (14.29)	24.33 (9.71)	−4
Caregiver Burden (higher scores = more burden)	20.25 (7.41)	15.5 (6.35)	−4.75	20.66 (8.08)	20.66 (7.76)	0.00

insurance. The majority of patients had Medi-Cal (Medicaid) or no insurance. While there are some rehabilitation options through counties that accept patients without insurance or Medi-Cal, they often have waitlists. Due to the brief intervention period, it was difficult to see rehabilitation outcomes. Relatedly, it is difficult to see outcomes related to employment since many patients are still recovering from the TBI during the intervention. A post-survey administered three months after completion of the intervention or control group will be used in the efficacy study to capture possible delayed effects of the intervention on employment.

Lastly, there were some important findings in the acceptability of the intervention. The intervention was designed to be administered within the first three months after patients experienced the TBI; however, patients and health professionals felt that this limited access for patients with TBI who often return home from the hospital, feel okay, and then realize months later that they are not progressing in their recovery and need support, yet are often cut off from healthcare resources by that time. Thus, eligibility criteria was expanded to one year or 365 days since hospitalization or rehabilitation for a TBI. Secondly, the original name of the intervention was *Promotora*, which is widely used in research literature.<sup>21</sup> However, many community members including patients, caregivers, and professionals, were not familiar with that word thus the name of the intervention was changed to *Trabajadora de Salud* since this name was familiar to community members and accurately described the intervention.

While the study's findings have implications for practice and future research, some limitations of the study include a small sample size, a post-test only administered at the end of the intervention with no follow-up, and a smaller sample size in the intervention group than control group. The small sample was used to pilot or test the acceptability and feasibility of the *Trabajadora de Salud* intervention prior to use of more resources to conduct an efficacy study. A follow-up post-test administered three or more months after the end of the intervention would be ideal, yet required more resources than were available to the author at the time of this pilot study. All participants were randomly assigned to the intervention or control groups with an equal chance of receiving the intervention or control group. Five were randomly selected into the control group while three were randomly selected into the intervention group using an application in Excel. It would improve future research on this topic to have equal sample sizes in the intervention and control groups.

## Conclusion

*Trabajadora de Salud* demonstrated promise in improving outcomes of Latinas/os with TBIs and their caregivers and should be further studied among larger sample sizes to assess its efficacy. The *Trabajadora de Salud* intervention should be considered as an option in improving the functional, depression, and somatic symptoms of the Latina/o patients with TBI as well as the somatic symptoms and caregiver burden of the caregivers.

## Funding

This study was funded by the Solution Focused Brief Therapy Association.

## References

- Haider AH, Chang DC, Efron DT, Haut ER, Crandall M, Cornwell EE. Race and insurance status as risk factors for trauma mortality. *Arch Surg*. 2008;143(10):945–949.
- Burnett DM, Silver TM, Kolakowsky-Hayner SA, Cifu DX. Functional outcome for African Americans and Hispanics treated at a traumatic brain injury model systems centre. *Brain Inj*. 2000;14(8):713–718.
- Haider AH, Chang DC, Efron DT, Haut ER, Crandall M, Cornwell EE. Race and insurance status as risk factors for trauma mortality. *Arch Surg*. 2008;143(10):945–949.
- Chang PFJ, Ostir GV, Kuo YF, Granger CV, Ottenbacher KJ. Ethnic differences in discharge destination among older patients with traumatic brain injury. *Arch Phys Med Rehabil*. 2008;89(2):231–236.
- Arango-Lasprilla JC, Ketchum JM, Gary KW, et al. The influence of minority status on job stability after traumatic brain injury. *Phys Med Rehabil*. 2009;1(1):41–49.
- Arango-Lasprilla JC, Ketchum JM, Drew A, et al. Neurobehavioural symptoms 1 year after traumatic brain injury: a preliminary study of the relationship between race/ethnicity and symptoms. *Brain Inj*. 2012;26(6):814–824.
- Gary KW, Arango-Lasprilla JC, Stevens LF. Do racial/ethnic differences exist in post-injury outcomes after TBI? A comprehensive review of the literature. *Brain Inj*. 2009;23(10):775–789.
- Armengol CG. A multimodal support group with Hispanic traumatic brain injury survivors. *J Head Trauma Rehabil*. 1999;14(3):233–246.
- Kreutzer JS, Stejskal TM, Ketchum JM, Marwitz JH, Taylor LA, Menzel JC. A preliminary investigation of the brain injury family intervention: impact on family members. *Brain Inj*. 2009;23(6):535–547.
- Perrin PB, Johnston A, Vogel B, et al. A culturally sensitive transition assistance program for stroke caregivers: examining caregiver mental health and stroke rehabilitation. *J Rehabil Res Dev*. 2010;47(7):605–617.
- Ayala GX, Vaz L, Earp JA, Elder JP, Cherrington A. Outcome effectiveness of the lay health advisor model among Latinos in the United States: an examination by role. *Health Educ Res*. 2010;25(5):815–840.
- Ayala GX, Vaz L, Earp JA, Elder JP, Cherrington A. Outcome effectiveness of the lay health advisor model among Latinos in the United States: an examination by role. *Health Educ Res*. 2010;25(5):815–840.
- Patient Protection and Affordable Care Act*, 42 U. S. C. 18001. 2010.
- Saverino C, Swaine B, Jaglal S, et al. Rehospitalization after traumatic brain injury: a population-based study. *Arch Phys Med Rehabil*. 2016;97(2):S19–S25.
- Stineman MG, Shea JA, Jette A, et al. The functional independence measure: tests of scaling assumptions, structure, and reliability across 20 diverse impairment categories. *Arch Phys Med Rehabil*. 1996;77(11):1101–1108.
- Seel RT, Kreutzer JS, Sander AM. Concordance of patients' and family members' ratings of neurobehavioral functioning after traumatic brain injury. *Arch Phys Med Rehabil*. 1997;78(11):1254–1259.
- Perrin PB, Johnston A, Vogel B, et al. A culturally sensitive transition assistance program for stroke caregivers: examining caregiver mental health and stroke rehabilitation. *J Rehabil Res Dev*. 2010;47(7):605–617.
- Perrin PB, Johnston A, Vogel B, et al. A culturally sensitive transition assistance program for stroke caregivers: examining caregiver mental health and stroke rehabilitation. *J Rehabil Res Dev*. 2010;47(7):605–617.
- Ayala GX, Vaz L, Earp JA, Elder JP, Cherrington A. Outcome effectiveness of the lay health advisor model among Latinos in the United States: an examination by role. *Health Educ Res*. 2010;25(5):815–840.
- Vincent D, Pasvogel A, Barrera L. A feasibility study of a culturally tailored diabetes intervention for Mexican Americans. *Biol Res Nurs*. 2007;9(2):130–141.
- Marsiglia FF, Bermudez-Parsai M, Coonrod D. Familias Sanas: an intervention designed to increase rates of postpartum visits among Latinas. *J Health Care Poor Underserved*. 2010;21(3 Suppl):119.